

On the Equivalence Principle of Quantum Mechanics

José M. Isidro

Dipartimento di Fisica “G. Galilei”
Istituto Nazionale di Fisica Nucleare
Università di Padova, Via Marzolo, 8
35131 Padova, Italy
`isidro@pd.infn.it`

February 1, 2008

Abstract

A recent concept in theoretical physics, motivated in string duality and M-theory, is the notion that not all quantum theories arise from quantising a classical system. Also, a given quantum model may possess more than just one classical limit. In view of these developments, we analyse some general properties that quantum mechanics must satisfy, if it is *not* to be formulated as a quantisation of a given classical mechanics. Instead, our approach to quantum mechanics is modelled on a statement that is close in spirit to the equivalence principle of general relativity, thus bearing a strong resemblance with the equivalence principle of quantum mechanics formulated by Faraggi–Matone.

1 Introduction

1.1 Motivation

Quantisation may be understood as a prescription to construct a quantum theory from a given classical theory. As such, it is far from being unique. Beyond canonical quantisation and Feynman’s path–integral, a number of different, often complementary approaches to quantisation are known, each one of them exploiting different aspects of the underlying classical theory. To mention just a few, the geometric quantisation of Kirillov, Kostant and Souriau relies heavily on the theory of group representations [1, 2, 3, 4]. Berezin’s quantisation can be applied to classical systems whose phase space is a homogeneous Kähler manifold [5]. The approach of Kontsevich [6] is based on the deformation quantisation of Poisson manifolds [7, 8], bearing some resemblance with noncommutative geometry.

The deep link existing between classical and quantum mechanics has of course been known for long. Perhaps its simplest manifestation is that of coherent states [9]. More recent is the notion that *not all quantum theories arise from quantising a classical system*. Furthermore, a given quantum model *may possess more than just one classical limit*. These ideas find strong evidence in string duality and M-theory [10, 11].

It therefore seems natural to try an approach to quantum mechanics that is not based, at least primarily, on the the quantisation of a given classical dynamics. In such an approach one would *not* take a classical theory as a starting point. Rather, quantum mechanics itself would be more fundamental, in that its classical limit or limits (possibly more than one) would follow from a parent quantum theory.

In order to carry out this programme one can envisage two different approaches to first quantisation, one technical, the other conceptual. A technical approach has been presented in [12], in connection with the quantum-mechanical implementation of an S-duality symmetry. In this paper we would like to take a more conceptual viewpoint instead.

1.2 Summary

The general purpose of this paper is to analyse some properties that quantum mechanics must satisfy, if it is *not* to be formulated as a quantisation of a given classical mechanics. We will formulate a statement, close in spirit to the equivalence principle of general relativity, that could well provide a starting point for a reformulation of quantum mechanics such as that claimed by Vafa [11]. Our formalism may be understood as a certain limit of Berezin's quantisation [5]. The latter relies on the metric properties of classical phase space $\mathcal{M}$, whenever $\mathcal{M}$ is a homogeneous Kähler manifold. In Berezin's method, quantum numbers arise naturally from the metric on $\mathcal{M}$. The semiclassical regime is then identified with the regime of large quantum numbers. Our method may be regarded as the topological limit of Berezin's quantisation, in that the metric dependence has been removed. As a consequence of this topological nature our quantum mechanics exhibits some added features. Quantum numbers are not originally present in our prescription; they appear only after a vacuum has been chosen, and even then they are local in nature, instead of global. Hence our procedure may be thought of as a manifestly non-perturbative formulation of quantum mechanics, in that we take no classical phase space and no Poisson brackets as our starting point, *i.e.*, we do not deform a classical theory into its quantum counterpart, as in deformation quantisation.

1.3 Outline

This paper is organised as follows. Section 2 sets the scene by giving a quick review of Berezin's quantisation, starting from the metric on certain classical phase spaces. Section 3 summarises the equivalence principle of Faraggi-Matone [13], as a preparation for our own starting point. The latter is presented under

the form of a statement in section 4. Its physical implications are analysed and discussed in detail in section 5.

2 Berezin's Quantisation

Below we sketch the construction of the Hilbert space of states from the metric on some relevant homogeneous Kähler manifolds [5].

2.1 The Complex Plane

Coherent states $|z\rangle$ of standard quantum mechanics on $\mathbf{R}$ are defined as the eigenstates of the annihilation operator $a = (Q + iP)/\sqrt{2\hbar}$. The Hilbert space of states is the Fock-Bargmann space $\mathcal{F}_\hbar(\mathbf{C})$ of entire analytic functions $\psi(z)$ on $\mathbf{C}$ with finite norm with respect to the scalar product

$$\langle\psi_1|\psi_2\rangle = \int_{\mathbf{C}} d\mu(z, \bar{z}) \exp(-z\bar{z}) \bar{\psi}_1(z) \psi_2(z), \quad (1)$$

the measure being given by $d\mu(z, \bar{z}) = \pi^{-1} dz \wedge d\bar{z}$. The latter follows from the Kähler form $\omega(z, \bar{z}) = dz \wedge d\bar{z}$ on $\mathbf{C}$. The argument of the exponential equals the Kähler potential $K_{\mathbf{C}}(z, \bar{z}) = z\bar{z}$ for the flat metric $g_{z\bar{z}} = 1$. There is a natural isomorphism between the oscillator states $|n\rangle$ and the basis states $z^n/\sqrt{n!}$, $n = 0, 1, \dots$ of the Fock-Bargmann space. The semiclassical limit corresponds to letting $n \rightarrow \infty$.

2.2 The Riemann Sphere

On the Riemann sphere S^2 , the Kähler potential $K_{S^2}(z, \bar{z}) = \log(1 + |z|^2)$ produces an integration measure $d\mu(z, \bar{z}) = (2\pi i)^{-1} dz \wedge d\bar{z}/(1 + |z|^2)^2$. The Hilbert space of states now becomes the space $\mathcal{F}_\hbar(S^2)$ of holomorphic functions on S^2 with finite norm, the scalar product being

$$\langle\psi_1|\psi_2\rangle = \left(\frac{1}{\hbar} + 1\right) \int_{S^2} d\mu(z, \bar{z}) (1 + |z|^2)^{-1/\hbar} \bar{\psi}_1(z) \psi_2(z). \quad (2)$$

It turns out that $\hbar^{-1}$ must be an integer. For ψ to have finite norm, it must be a polynomial of degree less than $\hbar^{-1}$. In fact, setting $\hbar^{-1} = 2j + 2$, $\mathcal{F}_\hbar(S^2)$ is the representation space for the spin- j representation of $SU(2)$. The semiclassical regime now corresponds to $j \rightarrow \infty$. Coherent states $|u\rangle$ are parametrised by points u in the quotient space $S^2 = SU(2)/U(1)$.

2.3 The Lobachevsky Plane

Consider the Lobachevsky plane modelled as the unit disc $D = \{z \in \mathbf{C} : |z| < 1\}$. From the Kähler potential $K_D(z, \bar{z}) = -\log(1 - |z|^2)$ one derives an integration measure $d\mu(z, \bar{z}) = (2\pi i)^{-1} dz \wedge d\bar{z}/(1 - |z|^2)^2$. With respect to the scalar

product

$$\langle \psi_1 | \psi_2 \rangle = \left(\frac{1}{\hbar} - 1 \right) \int_D d\mu(z, \bar{z}) (1 - |z|^2)^{1/\hbar} \bar{\psi}_1(z) \psi_2(z), \quad (3)$$

the space $\mathcal{F}_\hbar(D)$ of analytic functions on D with finite norm defines a Hilbert space of states. Setting $k - 1 = (2\hbar)^{-1}$, $\mathcal{F}_\hbar(D)$ becomes the representation space for the discrete series of the group $SU(1, 1)$ of isometries of D , *i.e.*, the space of weight- k modular forms. Large values of k correspond to the semiclassical limit of this quantum mechanics. Coherent states $|w\rangle$ are parametrised by points w in the quotient space $SU(1, 1)/U(1)$.

2.4 Complex Homogeneous Kähler Manifolds

Let $z^j, \bar{z}^k, j, k = 1, \dots, n$, be local coordinates on a complex homogenous Kähler manifold $\mathcal{M}$, and let $K_{\mathcal{M}}(z^j, \bar{z}^k)$ be a Kähler potential for the metric $ds^2 = g_{j\bar{k}} dz^j d\bar{z}^k$. The Kähler form $\omega = g_{j\bar{k}} dz^j \wedge d\bar{z}^k$ gives rise to an integration measure $d\mu(z, \bar{z})$,

$$d\mu(z, \bar{z}) = \omega^n = \det(g_{j\bar{k}}) \prod_{l=1}^n \frac{dz^l \wedge d\bar{z}^l}{2\pi i}. \quad (4)$$

The Hilbert space of states is the space $\mathcal{F}_\hbar(\mathcal{M})$ of analytic functions on $\mathcal{M}$ with finite norm, the scalar product being

$$\langle \psi_1 | \psi_2 \rangle = c(\hbar) \int_{\mathcal{M}} d\mu(z, \bar{z}) \exp(-\hbar^{-1} K_{\mathcal{M}}(z, \bar{z})) \bar{\psi}_1(z) \psi_2(z), \quad (5)$$

and $c(\hbar)$ a normalisation factor. Let G denote the Lie group of motions of $\mathcal{M}$, and assume $K_{\mathcal{M}}(z, \bar{z})$ is invariant under G . Setting $\hbar = k^{-1}$, the family of Hilbert spaces $\mathcal{F}_\hbar(\mathcal{M})$ provides a discrete series of projectively unitary representations of G . The homogeneity of $\mathcal{M}$ is used to prove that the correspondence principle is satisfied in the limit $k \rightarrow \infty$. Furthermore, let $G' \subset G$ be a maximal isotropy subgroup of the vacuum state $|0\rangle$. Then coherent states $|\zeta\rangle$ are parametrised by points ζ in the coset space G/G' .

3 The Equivalence Principle of Faraggi–Matone

In [13] an entirely new presentation of quantum mechanics has been given, starting from the so-called *equivalence principle of quantum mechanics*. In plain words, the philosophy underlying this approach could be summarised as follows. The classical Hamilton–Jacobi technique is based on transforming an arbitrary dynamical system, by means of coordinate changes, into a freely-evolving system subject to no interactions. The requirement that this equivalence also hold in the case when the conjugate variables are considered as dependent leads to a quantum analogue of the Hamilton–Jacobi equation corresponding to the one previously assumed by Floyd [14]. See also the book by Carroll [15].

A surprising new feature of the Faraggi–Matone approach is that the quantum analogue of the Hamiltonian characteristic function is quite different from the usual one in the literature. This solves Einstein’s criticism to Bohm’s approach (see [16] p. 243). As a result, the quantum potential is never trivial. The latter has been used in [17] to derive the gravitational interaction. This suggests that gravitation would have a quantum origin, so that physical interactions would *not* have to be introduced by hand, through the consideration of a potential. Rather, they would *follow* from consistency requirements. In particular, the quantum potential term that it is customary to neglect in the semiclassical limit plays a decisive role in determining the interaction, precisely due to the observation that it should not be neglected.

4 The Opening Statement

The equivalence principle of general relativity states that [18]

At every spacetime point in an arbitrary gravitational field, it is possible to choose a locally inertial coordinate system such that, within a sufficiently small region of the point in question, the laws of motion take the same form as in unaccelerated Cartesian coordinate systems in the absence of gravitation.

Let us now make the opening statement that

Given any quantum system, there always exists a coordinate transformation that transforms the system into the semiclassical regime, i.e., into a system that can be studied by means of a perturbation series in powers of $\hbar$ around a certain local vacuum.

One can perceive a conceptual analogy with the equivalence principle of Faraggi–Matone in the use of coordinate transformations in order to trivialise a given system. In our context, however, *trivialisation* does not mean cancellation of the interaction term, as in the Hamilton–Jacobi approach. Rather, it refers to the choice of a vacuum around which to perform a perturbative expansion in powers of $\hbar$. As we will see presently, this is equivalent to eliminating the metric, thus rendering quantum mechanics metrically trivial.

5 Physical Discussion

5.1 The Choice of a Vacuum

The opening statement above instructs us to choose a local vacuum. Under the *choice of a vacuum* we understand a specific set of coordinates around which to perform an expansion in powers of $\hbar$. This choice of a vacuum is *local in nature*, in that it is linked to a specific choice of coordinates. It breaks the group of allowed coordinate transformations to a (possibly discrete) subgroup, leaving behind a (possibly discrete) duality symmetry of the quantum theory. Call q the local coordinate corresponding to the vacuum in question, and Q its quantum operator. The corresponding local momentum P satisfies the usual Heisenberg algebra with Q . However, as our starting point we have no classical phase space

at all, and no Poisson brackets to quantise into commutators. This may be regarded as a manifestly non-perturbative formulation of quantum mechanics, such as that claimed in [11].

5.2 Quantum Numbers vs. a Topological Quantum Mechanics

Berezin's quantisation relied heavily on the metric properties of classical phase space. The semiclassical limit could be defined as the regime of large quantum numbers. The very existence of quantum numbers was a consequence of the metric structure. If quantum mechanics is *not* to be formulated as a quantisation of a given classical mechanics, then we had better do away with global quantum numbers, *i.e.*, with the metric. Metric-free theories usually go by the name of topological theories. Hence our quantum mechanics will be a *topological quantum mechanics*, *i.e.* free of *global* quantum numbers. Locally, of course, quantum numbers do appear, but only after the choice of a *local* vacuum.

5.3 Classical vs. Quantum

A feature of this approach is the following. After the choice of a local vacuum to expand around, the local quantum numbers one obtains describe excitations around the local vacuum chosen. Hence what appears to be a semiclassical excitation to a local observer may well turn out to be a highly quantum phenomenon, when described from the viewpoint of a different local vacuum.

The logic could be summarised as follows:

1. the fact that this quantum mechanics is topological implies the absence of a metric;
2. the absence of a metric implies the absence of global quantum numbers;
3. the absence of global quantum numbers implies the impossibility of globally defining a semiclassical regime. The latter exists only locally.

5.4 An example

In order to explore the quantum dynamics of a given system we need a knowledge of the Hilbert space of states and of the operators acting on it. Our opening statement does not determine them, so they can only be specified on a case-by-case basis, by imposing consistency with the symmetry requirements of the particular problem at hand. The same applies to the group of coordinate transformations that our opening statement refers to. We illustrate these points with the particular case of a point particle on the real line analysed in [12].

The basic requirement imposed in [12] is that an S-duality transformation (modelled on $\mathbf{Z}_2$) exist between the semiclassical and the strong quantum regimes. This determines the group of coordinate transformations to be $\mathrm{SL}(2, \mathbf{R})$. The Hilbert space of a point particle on the real line turns out to be

$L^2(0, \infty)$, which strictly contains $L^2(\mathbf{R})$. States belonging to $L^2(0, \infty)$ but not to $L^2(\mathbf{R})$ may be interpreted as nonperturbative states, that cannot be reached by the standard perturbative approach. The usual Hilbert space $L^2(\mathbf{R})$ only emerges after the choice of a vacuum. There are two possible vacua, $|0_z\rangle$ and $|0_{\bar{z}}\rangle$, corresponding to the coordinates z and $\bar{z} = -z^{-1}$ on the upper half-plane $\mathbf{H}$. The $\mathrm{SL}(2, \mathbf{R})$ symmetry acting homogeneously on $\mathbf{H}$ breaks down to the affine group, times $\mathbf{Z}_2$. The latter implements the desired S-duality symmetry exchanging the semiclassical and the strong quantum regimes of one single quantum theory. The coherent states constructed around $|0_z\rangle$ are not coherent from the viewpoint of $|0_{\bar{z}}\rangle$. This model does not allow for globally defined coherent states such as those of section 2, thus providing an explicit example of the general procedure presented above. Thus the quantum mechanics developed in [12] is that of affine variables on the real line. (Incidentally, affine symmetry is well-suited to quantum gravity [19, 20, 21, 22]).

Acknowledgments

It is a very great pleasure to thank Marco Matone for many conversations that made the preparation of this work much more enjoyable. The author would also like to thank Diego Bellisai, Kurt Lechner, Pieralberto Marchetti, Paolo Pasti, Dmitri Sorokin and Mario Tonin. This work has been supported by a Fellowship from Istituto Nazionale di Fisica Nucleare (Italy).

References

- [1] A. Kirillov, *Elements of the Theory of Representations*, Springer-Verlag, Berlin (1976).
- [2] B. Kostant, *Lectures in Modern Analysis and Applications III*, ed. by C. Taam, Lecture Notes in Mathematics, vol. 170, Springer-Verlag, Berlin (1970); *Group Representations in Mathematics and Physics*, ed. by V. Bargmann, Lecture Notes in Physics, vol. 6, Springer-Verlag, Berlin (1970).
- [3] J.-M. Souriau, *Group Theoretical Methods in Physics*, ed. by A. Janner, T. Janssen and M. Boon, Lecture Notes in Physics, vol. 50, Springer-Verlag, Berlin (1976).
- [4] N. Woodhouse, *Geometric Quantization*, Oxford University Press, Oxford (1991).
- [5] F. Berezin, *Sov. Math. Izv.* **38** (1974) 1116; *Sov. Math. Izv.* **39** (1975) 363; *Comm. Math. Phys.* **40** (1975) 153; *Comm. Math. Phys.* **63** (1978) 131.
- [6] M. Kontsevich, [q-alg/9709040](#); *Lett. Math. Phys.* **48** (1999) 35.
- [7] A. Cattaneo, G. Felder and L. Tomassini, [math.QA/0012228](#); G. Felder and B. Shoikhet, [math.QA/0002057](#). A. Cattaneo and G. Felder, [math.QA/0102108](#); [hep-th/0102208](#).

- [8] M. Schlichenmaier, [math.QA/0005288](#); [math.QA/0009219](#); A. Karabegov and M. Schlichenmaier, [math.QA/0006063](#); [math.QA/0102169](#).
- [9] J. Klauder and B.-S. Skagerstam, *Coherent States*, World Scientific, Singapore (1985); A. Perelomov, *Generalized Coherent States and their Applications*, Springer Texts and Monographs in Physics, Berlin (1986).
- [10] J. Schwarz, *Nucl. Phys. Proc. Suppl.* **55B** (1997) 1.
- [11] C. Vafa, [hep-th/9702201](#).
- [12] J.M. Isidro, [hep-th/0009221](#), to appear in *Int. Journ. Mod. Phys. A*.
- [13] A. Faraggi and M. Matone, *Phys. Lett.* **A249** (1998) 180; *Phys. Lett.* **B437** (1998) 369; *Phys. Lett.* **B445** (1998) 77; *Phys. Lett.* **B445** (1998) 357; *Phys. Lett.* **B450** (1999) 34; *Int. Journ. Mod. Phys. A***15** (2000) 1869; G. Bertoldi, A. Faraggi and M. Matone, *Class. Quant. Grav.* **17** (2000) 3965.
- [14] E. Floyd, *Found. Phys. Lett.* **9** (1996) 489; *Found. Phys. Lett.* **13** (2000) 235; *Int. Journ. Mod. Phys. A***14** (1999) 1111; *Int. Journ. Mod. Phys. A***15** (2000) 1363; [quant-ph/0009070](#).
- [15] R. Carroll, “Quantum Theory, Deformation and Integrability”, North-Holland Mathematics Studies **186**, Elsevier, Amsterdam (2000).
- [16] P. Holland, “The Quantum Theory of Motion”, Cambridge University Press, Cambridge (1993).
- [17] M. Matone, [hep-th/0005274](#).
- [18] S. Weinberg, “Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity”, Wiley, New York (1972).
- [19] C. Isham and A. Kakas, *Class. Quant. Grav.* **1** (1984) 621; *Class. Quant. Grav.* **1** (1984) 633.
- [20] J. Klauder, in *Relativity*, eds. M. Carmeli, S. Fickler and L. Witten, Plenum Press (1970).
- [21] J. Klauder and E. Alaksen, *Phys. Rev.* **D2** (1970) 272. I. Daubechies, J. Klauder and T. Paul, *Journ. Math. Phys.* **28** (1987) 85.
- [22] G. Watson and J. Klauder, *Journ. Math. Phys.* **41** (2000) 8072.